



Life in hot acid: pathway analyses in extremely thermoacidophilic archaea

Kathryne S Auernik, Charlotte R Cooper and Robert M Kelly

The extremely thermoacidophilic archaea are a particularly intriguing group of microorganisms that must simultaneously cope with biologically extreme pHs (≤ 4) and temperatures ($T_{\text{opt}} \geq 60$ °C) in their natural environments. Their expanding biotechnological significance relates to their role in biomining of base and precious metals and their unique mechanisms of survival in hot acid, at both the cellular and biomolecular levels. Recent developments, such as advances in understanding of heavy metal tolerance mechanisms, implementation of a genetic system, and discovery of a new carbon fixation pathway, have been facilitated by the availability of genome sequence data and molecular genetic systems. As a result, new insights into the metabolic pathways and physiological features that define extreme thermoacidophily have been obtained, in some cases suggesting prospects for biotechnological opportunities.

Address

Department of Chemical and Biomolecular Engineering, North Carolina State University, Raleigh, NC 27695-7905, United States

Corresponding author: Kelly, Robert M (rmkelly@eos.ncsu.edu)

Current Opinion in Biotechnology 2008, **19**:445–453

This review comes from a themed issue on
Tissue, Cell and Pathway Engineering
Edited by William Bentley and Michael Betenbaugh

Available online 11th September 2008

0958-1669/\$ – see front matter
© 2008 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.copbio.2008.08.001](https://doi.org/10.1016/j.copbio.2008.08.001)

Introduction

Over the past 20 years much has been written about the biotechnological potential of microorganisms from extreme environments, primarily focusing on individual enzymes capable of withstanding the otherwise harsh conditions required for long-term efficacy in bioprocessing environments [1–5]. However, as genome sequence data have become available for extremophiles (The UCSC Archaeal Genome Browser; <http://archaea.ucsc.edu>) and molecular genetics tools have begun to emerge [6,7], there exists the possibility to go beyond single biocatalytic steps to take advantage of the novel pathways and physiological characteristics that are intrinsic to these unique microorganisms. By incorporating these features into less extreme organisms and cells and by metaboli-

cally engineering extremophiles directly, a new horizon in microbial biotechnology can emerge. Here, we consider the extremely thermoacidophilic archaea, microorganisms that thrive in hot acid.


Extremely thermoacidophilic archaea and their physiological characteristics

For the purposes of this review, an ‘extreme thermoacidophile’ is a microorganism with both an optimal growth temperature ≥ 60 °C and an optimal pH of ≤ 4.0 . A majority of the extremely thermoacidophilic species studied to date [8] belongs to the archaeal orders of Sulfolobales and Thermoplasmatales (Figure 1). From what is currently known, it is interesting that the most heat-tolerant extreme thermoacidophiles are not the most acid-tolerant and vice versa. The most thermophilic of the extreme thermoacidophiles, crenarchaeon *Acidianus infernus*, grows at temperatures up to 95 °C (T_{opt} of 85–90 °C) but at pHs only as low as 1.0 (pH_{opt} 2.0) [9]. By contrast, *Picrophilus* species of the euryarchaeal order Thermoplasmatales are the most acidophilic, growing at pHs as low as 0 (pH_{opt} 0.7), but at temperatures up to only 65 °C (T_{opt} of 60 °C) [10]. Insights into life in hot acid may be forthcoming, since genome sequences exist or are underway for many extreme thermoacidophiles (see Figure 1 and Table 1) (www.genomesonline.org). Furthermore, several new species in known genera of Sulfolobales (*Acidianus*, *Metallosphaera*) have been reported, as well as a new member of the Thermoplasmatales, *Thermogymnomonas acidicola* [11–14]. We may have only scratched the surface with respect to extreme thermoacidophile diversity, because new environments not previously known to harbor these microorganisms have recently been identified. For example, mathematical modeling and 16S rRNA data suggest that conditions conducive to thermoacidophilic growth exist in deep-sea hydrothermal vents, a hypothesis supported by reports of the first euryarchaeon from the order Deep-sea Hydrothermal Vent Euryarchaeotic 2 (DHVE2), *Aciduliprofundum boonei* [15•]. Although *A. boonei* (T_{opt} 70 °C) grows best at a pH slightly above 4.0, 16S rRNA indicates that this microorganism comprises 10–15% of selected vent-associated archaeal populations, suggesting that there are extreme thermoacidophiles from these sites yet to be isolated [15•].

Mechanisms of resistance to and survival in hot acid

The mechanisms by which microbial life thrives in hot acid have been investigated in some detail in recent years,

Figure 1



Unrooted 16S phylogenetic tree (constructed using ClustalW and Phylip 3.5c available on <http://mobyli.pasteur.fr>) of extremolythermoacidophilic archaea ($T_{opt} \geq 60$ °C, $pH_{opt} < 4$), compiled from [8,9,11,14,72,73]. Genomes are denoted as ^Ssequenced or ^{Sip}sequencing in progress, according to www.genomesonline.org. Organisms denoted in quotes have not yet been described in full detail. Accession numbers for 16S rRNA are listed in parentheses. Sso, *S. solfataricus* (SSO03); 'Sis, *S. islandicus*' strain M14A (AY247895); Ssh, *S. shibatae* (M32504); Sto, *S. tokodaii* (ABO22438); 'Syang, *Sulfolobus yangmingensis*' (ABO10957); Sowi, *Sulfurisphaera ohwakuensis* (D85507); Saci, *S. acidocaldarius* (D14876); 'Sthur, *Sulfolobus thuringensis*' (X90485); Sazo, *Stygiolobus azoricus* (D85520); Smet, *S. metallicus* (U40813); Mscd, *M. sedula* (Msed_R0026); Mpru, *Metallosphaera prunae* (X90482); Mhak, *Metallosphaera hakonensis* (D86414); Abr, *Acidianus brierleyi* (X90477); 'Atn, *Acidianus tengchongensis*' (AF226987); Ainf, *A. infernus* (X89852); Aam, *Acidianus ambivalens* (D85506); Asuf, *Acidianus sulfidivorans* (AY907891); Aac, *Acidolobus aceticus* (AF191225); Cmaq, *Caldivirga maquilingensis* (ABO13926); Posh, *Picrophilus oshimae* (X84901); Pto, *Picrophilus torridus* (PTO02); Taci, *Thermogymnomonas acidicola* (AB269873); Tvn, *Thermoplasma volcanium* (Tvn04); DHVE2 represented by 16S of *A. boonei* (DQ451875). No 16S sequences are available for extreme thermoacidophiles *Sulfurococcus yellowsonensis* and *Sulfurococcus mirabilis*.

triggered by the availability of genome sequence data, functional genomics tools, and molecular genetics. Although the intrinsic basis for this novel growth physiology is not clear, clues are emerging as to how these microorganisms survive in the face of hot, acidic, and often metal-laden conditions which are typically associated with DNA damage, protein denaturation, and other disruptions in cellular processes.

DNA damage and repair

High temperatures and the potential for cytosol acidification heighten the possibility of DNA damage or modification in extreme thermoacidophiles relative to mesophilic neutrophiles. Thus, clues to DNA damage repair may emerge from examination of this cellular function in hot acid biotopes. It is surprising that basal mutation rates for extreme thermoacidophiles are not particularly high. For example, *Sulfolobus acidocaldarius* has a spontaneous mutation rate similar to that of *E. coli* [16]. Furthermore, when *Sulfolobus solfataricus* and *S. acidocaldarius* were exposed to UV-irradiation, no significant increase in transcription of known DNA repair proteins was noted [17,18]; however, it is possible that these genes are constitutively transcribed at higher levels than in mesophiles. Following irradiation, aggregates resembling those formed during plasmid-mediated conjugation were found, spurring speculation that *Sulfolobus*

species may use conjugational DNA exchange and homologous recombination to repair mutated DNA [17]. In a related study, *S. solfataricus* infected with the *Sulfolobus* spindle-shaped virus (SSV1) exhibited a similar, but heightened, response to UV-induced DNA damage, suggesting that viruses may be an evolutionary component of stress management systems [19]. Another spindle-shaped virus, SSV2 from native host '*Sulfolobus islandicus*' REY15/4, sent infected '*S. islandicus*' REY15A cells into a metabolically inactive state upon encountering unfavorable environmental conditions and then played a role in restarting metabolic activity once favorable growth conditions emerged [20]. Up-regulation of two *S. solfataricus* *recA/rad51* homologs (*radA*, SSO0250; *radA*-like, SSO0777) in response to a DNA-damaging antibiotic led to the discovery of the first regulatory protein involved in archaeal DNA damage repair (Sta1, SSO0048) [21]. Neither the *radA*-like SSO0777, its *sta1* activator, nor *radA* itself was induced by UV-irradiation, indicating that the nature of DNA damage may drive the specific type of repair response [21].

Heat shock

Though extreme thermoacidophiles thrive at temperatures up to 95 °C, they are still susceptible to thermal stresses such that they exhibit both cold shock and heat shock responses. Extremely thermoacidophilic archaea

Table 1**Selected features of extreme thermoacidophiles with sequenced genomes**

Sequenced thermoacidophile	Growth temperature (°C)	Growth pH	Growth optimum (°C, pH)	Isolated from	Genome sequence reported	GC content (%)	Genome size (Mb)	Estimated # of encoded proteins	IS elements ^a	Toxin-like COGs ^b	Hypothetical proteins
Crenarchaea-Thermoprotei											
Thermoproteales											
<i>Caldivirga maquilingensis</i>	60–92	2.3–6.4	85, 3.7–4.2	Acidic hot spring, Philippines	2007	43.1	2.1	1963	7	15	549
Sulfolobales											
<i>Metallosphaera sedula</i>	50–80	1–4.5	75, 2–3	Solfataric thermal pond drainage, Italy	2007	46.2	2.2	2256	11	18	673
<i>Sulfolobus acidocaldarius</i>	55–85	1–6	70–75, 2–3	Solfataric hot spring, Italy	2005	36.7	2.2	2223	14	19	1002
<i>Sulfolobus solfataricus</i>	50–87	2–5.5	85, 3–4.5	Solfataric hot spring, Italy	2001	35.8	3	2977	159	26	1340
<i>Sulfolobus tokodaii</i>	70–85	2–5.5	80, 2.5–3	Hot spring, Japan	2001	32.8	2.7	2825	12	32	1874
Euryarchaea-Thermoplasmata											
Thermoplasmatales											
<i>Picrophilus torridus</i>	47–65	0–3.5	60, 0.7	Volcanic solfataric field, Japan	2004	36	1.5	1535	4	4	291
<i>Thermoplasma acidophilum</i>	45–63	0.5–4	59, 1–2	Burning coal refuse pile, USA	2001	46	1.6	1482	4	6	583
<i>Thermoplasma volcanium</i>	33–67	1–4	60, 2	Volcanic solfataric field, Italy	2001	39.9	1.6	1499	56	6	261

Compiled from NCBI May 2008 genome project links (<http://www.ncbi.nlm.nih.gov/>) and references listed for this article.

^a Annotation contains 'transposase', 'integrase', or 'resolvase'.


^b ORFs similar to COGs 1412, 1439, 1487, 1848, 3413, or 4113 containing nucleic acid binding and/or PIN (PiIT N terminus) domains.

react to supraoptimal temperatures in much the same way as other microorganisms [22–24]. Most work to date has focused on the archaeal thermosome, or rosettasome, a heat-shock responsive HSP60-like molecular chaperone that has been implicated in many cellular roles [25]. However, recent efforts have shown that heat shock response in extreme thermoacidophiles is extensive, involving much more than chaperones or other proteins involved in protein refolding. When exponentially growing *S. solfataricus* was shifted from its growth temperature optimum (80 °C) to 90 °C, approximately 1/3 of the transcriptome responded within 5 min [26]. Included in this set of genes were many insertion elements and chromosomally encoded toxin–antitoxin (TA) loci — 22 TA pairs and 1 solitary toxin — all from the VapBC family [26]. Chromosomally encoded TA loci in bacteria are thought to be stress response elements [27], though the role of these tandem protein complexes in archaea has not been examined. Since PIN domain-containing VapCs, the ‘toxin’ component of TA loci, are putative ribonucleases [28,29], these proteins could play an important role in post-transcriptional regulation in archaea, especially during heat shock.

Metal resistance

Extreme thermoacidophiles have developed mechanisms for tolerating heavy metals that are physiologically toxic to most microorganisms (Figure 2). These mechanisms involve their capacity to recover from metal-induced damage (similar to oxidative stress) [30] and to limit the effective concentration of the toxic metal itself. In some cases, enzymes reduce or oxidize metals to less toxic forms — for example, the mercuric reductase in *S. solfataricus* reduces soluble intracellular Hg^{2+} to volatile elemental Hg^0 [31•]. In other cases, metal chelation or complexation can accomplish the same objective. In *Sulfolobus metallicus*, a polyphosphate (polyP)-based mechanism is believed to underlie cellular tolerance to high levels of copper; greater accumulation of polyP granules was observed in *S. metallicus* (considered to have higher levels of Cu tolerance) compared to *S. solfataricus*, and granule size was noted to decrease as Cu levels were increased [32•]. Other strategies, however, do not involve metal transformation, direct or indirect, and instead are based on exporting toxic metal ions via P-type ATPases [33]. Evidence to date suggests that multiple systems can operate in parallel in extreme thermoacidophiles to provide cumulative tolerance, with particular strategies useful for multiple metals [31•,32•,33]. For example, copper tolerance in *Sulfolobus* species involves efflux ATPases in addition to the polyP pathway, but the same ATPases also contribute to cadmium tolerance [32•,33]. There may also be some intrinsic redundancy in protecting against heavy metal toxicity. Disruption mutants lacking mercuric reductase and its regulator (*merAR*) were found to still exhibit some *mer* operon transcription [31•]. In fact, creation of a mutant with a disrupted regulator resulted

Figure 2



Metal resistance mechanisms in extreme thermoacidophiles. CopA is the P-type ATPase shown to be involved in copper and cadmium cation efflux in *S. solfataricus* [33]. MerA is the mercuric reductase which reduces soluble Hg^{2+} to volatile elemental Hg and is constitutively expressed in *S. solfataricus* [31•]. Ppk (polyphosphate kinase) and Ppx (exopolyphosphatase) which comprise the polyP system described in *S. metallicus* [32•].

in increased *merA* expression and consequently increased Hg^{2+} tolerance [31•], underscoring the importance of understanding the regulation of resistance mechanisms with respect to engineering characteristics for bioprocesses. Although traditional acclimation and/or spontaneous mutant generation approaches are still useful, direct genetic manipulation offers the possibility of conferring similar levels of metal tolerance increase in a systematic manner.

Molecular genetics of extreme thermoacidophiles

Versatile genetic systems for extreme thermoacidophiles are a crucial need for many reasons. Recombinant expression of genes encoding extreme thermoacidophile proteins in commonly used bacterial hosts can be problematic [34,35], probably reflecting intrinsic differences between archaea and bacteria. Also, molecular genetic systems could provide the basis for investigating biological mechanisms enabling life in hot acid. Fortunately, promising developments along these lines have been reported, including successful efforts with gene disruption [36,37] and protein tagging [6]. For a comprehensive review of the development of molecular genetics for archaea see [7].

Viruses and plasmids

Virus-based plasmids have been tailored for specific needs to support the development of genetics systems in extreme thermoacidophiles. The first virus from an extreme thermoacidophile, SSV1, was isolated over 20

years ago from *Sulfolobus shibatae* (B12) [38] and is the one most developed for use in studying *Sulfolobus* species [7]. There are some useful features of this virus for molecular genetics. For example, the removal of the integrase gene from SSV1 demonstrated that the integrase protein is essential only for the integration into the host genome, and not for virus infection and replication [39]. This characteristic could be exploited for novel vector development, where nonhomologous recombination of the vector into the host genome is undesirable. Recently constructed *Sulfolobus*–*E. coli* fusion vectors have potential as easily modifiable genetic elements for extreme thermoacidophiles. The fusion shuttle vector pSSVrt, a combination of '*S. islandicus*' REY15/4 pSSVx and *E. coli* pUC19, accommodated insertion of foreign sequences up to ~11 kb with efficient propagation and vector stability at high-copy numbers with no integration [40]. Furthermore, the pRN1-*E. coli* transposon fusion, noted for its relatively small size (5.4 kb) and stable copy numbers of 10–20 in mid-log phase, is stable in *S. acidocaldarius* and *S. solfataricus*, as well as in *E. coli*. [41]. By adding the *pyrE* gene to this plasmid and using *pyrE* mutants, a selectable marker could be introduced into the host. An SSV1–*E. coli* pUC18 reporter gene system was developed with selectable marker genes *pyrEF*, both heat-inducible and arabinose-inducible promoters, and convenient restriction sites [42]. The improved vector (modified from pMJ03) was used for heterologous and homologous production of tagged proteins in *S. solfataricus*. All of these developments are exciting and offer promise for expanding molecular genetics capabilities in extreme thermoacidophiles.

Gene disruption

Construction of directed gene deletion mutants in extremely thermoacidophilic archaea is a significant challenge, but progress is being made on this front. The focus has been on *S. solfataricus* PBL2025, a constructed mutant of *S. solfataricus* 98/2, which lacks about 50 genes, including *lacS* [43]. Thus, the inability to grow on lactose-based minimal media provides a selectable marker. A protocol for efficient integration of exogenous DNA into the *S. solfataricus* PBL2025 genome has been described [36]. A similar method was used to construct a deficient mutant to study α -amylase function and regulation in *S. solfataricus* [37]. Markerless exchange using a plasmid that encodes a cloned copy of a modified DNA sequence and a selectable marker gene, again *lacS* for natural deletion mutant PBL2025, has also been used for the development of mercury reductase (*merA*)-deficient *Sulfolobus solfataricus* [31••].

Recombinant production of extremely thermoacidophilic proteins

Production of extremely thermoacidophilic proteins in mesophilic hosts (e.g. *E. coli*) can take the advantage of overexpression and simplified purification methods (e.g. heat treatment), but codon usage and inclusion body

problems often temper the enthusiasm for this approach [34,35]. Some solutions to existing problems have been recently proposed. *S. solfataricus* genes have rare (compared to *E. coli*) codon clustering at the 5' transcript end which specifically inhibits target translation. But, this can be relieved by adding rare codon tRNAs (utilizing strains like BL21(DE3) CodonPlus-RIL or Rosetta(DE3)) or by changing rare codons (via primer design) to those more frequently translated by the host [34]. Yields of active *S. tokodaii* and *P. torridus* proteins produced in *E. coli* (Rosetta(DE3)) were increased by growth and expression at elevated temperatures up to 46 °C, where it was hypothesized that protein synthesis was slowed, contributing to an increase in rate of proper folding [35].

Bioleaching

The biomining industry has a longstanding interest in the use of extreme thermoacidophiles for metals recovery from ores [44,45]. These organisms, as is the case with certain mesophilic chemolithotrophic bacteria such as *Acidithiobacillus ferrooxidans* [46•], can liberate precious (e.g. gold) and base (e.g. copper) metals trapped in, and as, metal sulfides (e.g. iron pyrite and chalcopyrite) through dissimilatory oxidative processes. Biological regeneration of Fe^{3+} from Fe^{2+} is the key to chemical attack of metal sulfides. However, biooxidation of reduced inorganic sulfur compounds (RISCs) is also important to prevent the accumulation of passivating sulfur compounds on metal surfaces that can limit metal mobilization rates. Extreme thermoacidophiles grow at temperatures where mesoacidophilic biocatalysts (or contaminants from nonsterile substrates) cannot, and where passivation from RISCs is nearly eliminated, lead to higher effective leaching rates [46•]. Efficacy in biomining environments also requires tolerance of high levels of toxic heavy metals as well as the ability to assimilate inorganic carbon, as organic sources can be scarce in this environment. A full complement of the aforementioned desirable traits is not typically resident in a single native microorganism, but may be in a consortium [46•]. Alternatively, extreme thermoacidophile genomes [47–49], including that of a biomining organism [50•], can be examined for pathways responsible for conferring these desirable biomining traits. Taken together with the emerging molecular genetic tools for extreme thermoacidophiles, metabolic engineering of biomining organisms with enhanced properties may soon be a reality.

Iron and sulfur oxidation

Several terminal components of respiratory electron transport chains (ETC) in extreme thermoacidophiles have been known for years [51–54]. However, it was recently shown that the relative membrane concentration of these components depends on the electron donating substrate [55], suggesting the involvement in dissimilatory Fe^{2+} and RISC oxidation. Substrate-dependent ETC expression led to the identification of a Fe^{2+} oxidation- (*fox*) induced gene

cluster in the autotrophic biominer, *S. metallicus* [56^{••}]. Comparative genomics subsequently revealed that homologs to this gene cluster are also present in *Metallosphaera sedula* [50[•]] and *Sulfolobus tokodaii*, the latter of which was not previously shown to have iron-oxidizing capabilities [56^{••}]. This cluster contains not only ORFs similar to previously recognized terminal components (*foxABCD*), but also ferredoxins and other putative iron-sulfur binding proteins typically involved in electron transfer (*foxFGHJ*). This implies that more than terminal components, possibly entire ETCs, are differentially expressed in response to certain substrates. Although the sulfur oxygenase reductase (SOR) does not appear to be directly connected to an ETC [57[•]], it was also up-regulated in S^0 compared to Fe^{2+} grown cells [56^{••}]. SOR is believed to participate in the early steps of cytoplasmic sulfur oxidation in extremely thermoacidophilic archaea. Surprisingly, *M. sedula*, a putative sulfur oxidizer [58[•]], does not encode SOR [50[•]], indicating that either some extremely thermoacidophilic archaea have alternative (yet unknown) sulfur oxidation enzymes, or that *M. sedula* has lost (or never possessed) the capacity to oxidize sulfur (similar to *S. solfataricus* and *S. acidocaldarius*) [58[•]].

CO₂ fixation


With the identification of a fifth pathway for inorganic carbon fixation [59^{••}], new perspectives on autotrophy have emerged. The new cycle model starts with a two-carbon acetyl-CoA molecule and assimilates two bicarbonates. The key bicarbonate-incorporating enzyme is a heterotrimeric acetyl-/propionyl-CoA carboxylase [60]. One way in which this new pathway is distinguished from the 3-hydroxypropionate cycle is by a putative homotetrameric 4-hydroxybutyryl-CoA dehydratase (Figure 3). Although activities have been detected for all steps in this pathway [59^{••}], only a few enzymes have been characterized biochemically [59^{••},60–62]. Comparative genomics suggests multiple candidates (all having similar sequences) for the final enzymatic reactions of the cycle; experimental work is required to identify which of these candidates supports the function of the final three steps of the pathway, involving rearrangement of a four-carbon molecule (crotonyl-CoA) before splitting into two molecules of acetyl-CoA. Understanding and controlling regulation of this new inorganic carbon fixation pathway could lead to the improvements in bioleaching through enhanced biomass/biocatalyst levels, and even new CO₂ sequestration strategies.

Other developments

ncRNA

Noncoding RNAs (ncRNAs) in extremely thermoacidophilic archaea, particularly in *Sulfolobus* species [63–66], could be used for transient control of biocatalytic steps in a bioprocess via dosing of small interfering RNA (siRNA). Although many ncRNAs are also small RNAs (sRNA) of 60 nucleotides or less, ncRNAs as long as 500 nucleotides

Figure 3



The recently proposed fifth cycle of autotrophic carbon fixation, adapted from [59^{••}]. (1) Acetyl-CoA/propionyl-CoA carboxylase, (2) malonyl-CoA/succinyl-CoA reductase, (3) malonate semialdehyde reductase, (4) 3-hydroxypropionyl-CoA synthetase, (5) 3-hydroxypropionyl-CoA dehydratase, (6) acryloyl-CoA reductase, (7) methylmalonyl-CoA epimerase, (8) methylmalonyl-CoA mutase, (9) succinate semialdehyde reductase, (10) 4-hydroxybutyryl-CoA synthetase, (11) 4-hydroxybutyryl-CoA dehydratase, (12) crotonyl-CoA hydratase, (13) 3-hydroxybutyryl-CoA dehydrogenase, (14) acetoacetyl-CoA β -ketothiolase. ORFs encoding enzymes with (?) have not been finalized in the extreme thermoacidophile in which the cycle was studied. Many of the intermediates of this cycle are the same as found in the 3-hydroxypropionate cycle (left side and base of triangle). The right side of triangle represents the rearrangement after the second carbon addition, concluding with a split into two molecules of acetyl-CoA, which distinguishes the 3-hydroxypropionate/4-hydroxybutyrate cycle from the 3-hydroxypropionate cycle.

have been reported [63]. Most ncRNAs are characterized by a K-turn motif and appear to possess a post-transcriptional modification or regulation (silencing) function. Many ncRNAs recognize their targets via full (antisense) or partial (antisense-box motif) complementarity to transposons, other ORFs, or noncoding RNA (rRNA, tRNA, sRNA, etc.) [63,64]. ncRNAs are thought to interact with proteins to influence structure and function. For example, in *S. solfataricus*, the Rbp18 protein of the 30S ribosomal subunit has been shown to bind free sRNA (*in vitro* and *in vivo*) with some degree of selectivity [65]. In *S. acidocaldarius*, the antisense-box motifs (and their location) in sRNA are important for both proper structural complexation with ribosomal core proteins and complex activity [64]. sRNA produced from clustered regularly interspaced short palindromic repeats (CRISPRs) are believed to interact with catalytically active CRISPR-associated sequences (Cas proteins), which are often encoded immediately upstream (crenarchaea) or downstream (euryarchaea) of CRISPRs [63,66]. Some CRISPRs contain nonrepetitive spacer sequences with similarity to viruses

or genomic ORFs. The Cas proteins may be involved with adding/removing spacer sequences to/from CRISPR regions (i.e. Cas1, Cas2), processing of long ncRNA to sRNA form (i.e. Cas3, Cas5), and possibly complexing with resulting sRNAs to form a microbial equivalent of the eukaryotic RNA-induced silencing (RISC) complex (i.e. Cas4) [66,67].

S-layers and extreme thermoacidophile membranes

S-layers of extremely thermoacidophilic archaea are of interest in nanobiotechnology, because their self-assembled periodicity and uniform properties, as well as thermoacid stability, make them attractive for the use in ultrafiltration, immobilization matrix, and coating capacities [68]. Most work to date with these S-layers has focused on structure and properties of the purified self-assembled proteins [69]. Models suggest that archaeal S-layers, with their membrane anchors, help maintain cell shape and stabilize the membrane against environment-induced osmotic pressure changes [70]. Study and use of tetraetherlipids themselves have been somewhat hampered due to high costs/low purification yields. However, recent efforts to defray bioleaching costs by processing by-products have resulted in a lower-cost, higher yield purification process for extremely thermoacidophilic tetraetherlipid, calditoglycero-caldarchaeol [71]. The next significant advances in functional understanding may come from studies of the natural environment of S-layers, in which they are bound to the tetraetherlipid cell membrane [69]. A search through GenBank (July 2008) shows that although S-layer domain proteins are annotated in sequenced extreme thermoacidophiles (COG 1361), genes related to their modification and assembly are not well known/annotated. Future work in this area will most probably begin with the analysis of gene neighborhoods (many containing transporters and/or transcriptional regulators). Manipulation via genetic tools will be invaluable in study of the formation and regulation of these proton influx barriers with long-term potential to produce tunable acid stability.

Summary

Recently available extreme thermoacidophile genome sequences are revealing novel pathways and strategies that contribute to the survival in hot, acidic environments. With the emerging availability of molecular genetics for these microorganisms, metabolic engineering efforts to realize biotechnological opportunities are within reach. Also promising is the prospect of finding novel extreme thermoacidophiles in yet untapped acidic niches, such as deep-sea hydrothermal biotopes.

Acknowledgements

KSA and CRC acknowledge NIH T32 Biotechnology Traineeships for support. This work was funded in part by a grant to RMK from the US National Science Foundation.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Adams MW, Perler FB, Kelly RM: **Extremozymes: expanding the limits of biocatalysis**. *Biotechnology (N Y)* 1995, **13**:662-668.
 2. Antranikian G, Vorgias CE, Bertoldo C: **Extreme environments as a resource for microorganisms and novel biocatalysts**. *Adv Biochem Eng Biotechnol* 2005, **96**:219-262.
 3. Hough DW, Danson MJ: **Extremozymes**. *Curr Opin Chem Biol* 1999, **3**:39-46.
 4. Tindall KR, Kunkel TA: **Fidelity of DNA synthesis by the *Thermus aquaticus* DNA polymerase**. *Biochemistry* 1988, **27**:6008-6013.
 5. Atomi H: **Recent progress towards the application of hyperthermophiles and their enzymes**. *Curr Opin Chem Biol* 2005, **9**:166-173.
 6. Sowers KR, DasSarma S, Blum PH: **Gene transfer in archaea**. In *Methods for General and Molecular Microbiology*. Edited by Reddy CA, Beveridge TJ, Breznak JA, Marzluf GA, Schmidt TM. American Society for Microbiology; 2007.
 7. Allers T, Mevarech M: **Archaeal genetics – the third way**. *Nat Rev Genet* 2005, **6**:58-73.
 8. Garrity GM, Lilburn TG, Cole JR, Harrison SH, Euzeby J, Tindall BJ: **Taxonomic Outline of the Bacteria and Archaea, Release 7.7**. Michigan State University Board of Trustees; 2007: 6–31.
 9. Huber H, Prangishvili D: **The Sulfolobales**. In *The Prokaryotes*, edn 3. Edited by Dworkin M, Falkow S, Rosenberg E, Schleifer K, Stackebrandt E. Springer; 2006:23-50.
 10. Huber H, Stetter KO: **Thermoplasmatales**. In *The Prokaryotes*, edn 3. Edited by Dworkin M, Falkow S, Rosenberg E, Schleifer K, Stackebrandt E. Springer; 2006:101-112.
 11. Plumb JJ, Haddad CM, Gibson JA, Franzmann PD: ***Acidianus sulfidivorans* sp. nov., an extremely acidophilic, thermophilic archaeon isolated from a solfatara on Lihir Island, Papua New Guinea, and emendation of the genus description**. *Int J Syst Evol Microbiol* 2007, **57**:1418-1423.
 12. Yoshida N, Nakasato M, Ohmura N, Ando A, Saiki H, Ishii M, Igarashi Y: ***Acidianus manzaensis* sp. nov., a novel thermoacidophilic archaeon growing autotrophically by the oxidation of H₂ with the reduction of Fe³⁺**. *Curr Microbiol* 2006, **53**:406-411.
 13. Kozubal M, Macur RE, Korf S, Taylor WP, Ackerman GG, Nagy A, Inskeep WP: **Isolation and distribution of a novel iron-oxidizing crenarchaeon from acidic geothermal springs in Yellowstone National Park**. *Appl Environ Microbiol* 2008, **74**:942-949.
 14. Itoh T, Yoshikawa N, Takashina T: ***Thermogymnomonas acidicola* gen. nov., sp. nov., a novel thermoacidophilic, cell wall-less archaeon in the order Thermoplasmatales, isolated from a solfataric soil in Hakone, Japan**. *Int J Syst Evol Microbiol* 2007, **57**:2557-2561.
 15. Reysenbach AL, Liu Y, Banta AB, Beveridge TJ, Kirshtein JD, •• Schouten S, Tivey MK, Von Damm KL, Voytek MA: **A ubiquitous thermoacidophilic archaeon from deep-sea hydrothermal vents**. *Nature* 2006, **442**:444-447.
- This study discusses strategy used to isolate a new order of extremely thermoacidophilic organisms in a low pH deep-sea environment.
16. Grogan DW, Carver GT, Drake JW: **Genetic fidelity under harsh conditions: analysis of spontaneous mutation in the thermoacidophilic archaeon *Sulfolobus acidocaldarius***. *Proc Natl Acad Sci U S A* 2001, **98**:7928-7933.
 17. Frols S, Gordon PM, Panlilio MA, Duggin IG, Bell SD, Sensen CW, • Schleper C: **Response of the hyperthermophilic archaeon *Sulfolobus solfataricus* to UV damage**. *J Bacteriol* 2007, **189**:8708-8718.
- Genome-wide transcriptional response analysis of Sso PH1 and Sso PH1 (SSV1) to UV irradiation to elucidate the DNA damage/repair mechanisms.

18. Gotz D, Paytubi S, Munro S, Lundgren M, Bernander R, White M: **Responses of hyperthermophilic crenarchaea to UV irradiation.** *Genome Biol* 2007, **8**:R220.
19. Frols S, Gordon PM, Panlilio MA, Schleper C, Sensen CW: **Elucidating the transcription cycle of the UV-inducible hyperthermophilic archaeal virus SSV1 by DNA microarrays.** *Virology* 2007, **365**:48-59.
20. Contursi P, Jensen S, Aucelli T, Rossi M, Bartolucci S, She Q: **Characterization of the *Sulfolobus* host-SSV2 virus interaction.** *Extremophiles* 2006, **10**:615-627.
21. Abella M, Rodriguez S, Paytubi S, Campoy S, White MF, Barbe J: **The *Sulfolobus solfataricus* radA paralogue sso0777 is DNA damage inducible and positively regulated by the Sta1 protein.** *Nucleic Acids Res* 2007, **35**:6788-6797.
22. Han CJ, Park SH, Kelly RM: **Acquired thermotolerance and stressed-phase growth of the extremely thermoacidophilic archaeon *Metallosphaera sedula* in continuous culture.** *Appl Environ Microbiol* 1997, **63**:2391-2396.
23. Trent JD, Osipiuk J, Pinkau T: **Acquired thermotolerance and heat shock in the extremely thermophilic archaeobacterium *Sulfolobus* sp. strain B12.** *J Bacteriol* 1990, **172**:1478-1484.
24. Peeples TL, Kelly RM: **Bioenergetic response of the extreme thermoacidophile *Metallosphaera sedula* to thermal and nutritional stresses.** *Appl Environ Microbiol* 1995, **61**:2314-2321.
25. Kagawa HK, Yaoi T, Brocchieri L, McMillan RA, Alton T, Trent JD: **The composition, structure and stability of a group II chaperonin are temperature regulated in a hyperthermophilic archaeon.** *Mol Microbiol* 2003, **48**:143-156.
26. Tachdjian S, Kelly RM: **Dynamic metabolic adjustments and genome plasticity are implicated in the heat shock response of the extremely thermoacidophilic archaeon *Sulfolobus solfataricus*.** *J Bacteriol* 2006, **188**:4553-4559.
27. Pandey DP, Gerdes K: **Toxin-antitoxin loci are highly abundant in free-living but lost from host-associated prokaryotes.** *Nucleic Acids Res* 2005, **33**:966-976.
28. Clissold PM, Ponting CP: **PIN domains in nonsense-mediated mRNA decay and RNAi.** *Curr Biol* 2000, **10**:R888-R890.
29. Daines DA, Wu MH, Yuan SY: **VapC-1 of nontypeable *Haemophilus influenzae* is a ribonuclease.** *J Bacteriol* 2007, **189**:5041-5048.
30. Salzano AM, Febbraio F, Farias T, Cetrangolo GP, Nucci R, Scaloni A, Manco G: **Redox stress proteins are involved in adaptation response of the hyperthermoacidophilic archaeon *Sulfolobus solfataricus* to nickel challenge.** *Microb Cell Fact* 2007, **6**:25.
31. Schelert J, Drozda M, Dixit V, Dillman A, Blum P: **Regulation of mercury resistance in the crenarchaeote *Sulfolobus solfataricus*.** *J Bacteriol* 2006, **188**:7141-7150.
- This study discusses the use of genetic tools in *S. solfataricus* to elucidate the regulation of a mercury resistance mechanism.
32. Remonsellez F, Orell A, Jerez CA: **Copper tolerance of the thermoacidophilic archaeon *Sulfolobus metallicus*: possible role of polyphosphate metabolism.** *Microbiology* 2006, **152**:59-66.
- A study of Cu efflux and correlation to polyP accumulation and enzyme activity in selected *Sulfolobus* species with high and low Cu tolerances.
33. Ettema TJ, Brinkman AB, Lamers PP, Kornet NG, de Vos WM, van der Oost J: **Molecular characterization of a conserved archaeal copper resistance (cop) gene cluster and its copper-responsive regulator in *Sulfolobus solfataricus* P2.** *Microbiology* 2006, **152**:1969-1979.
34. Kim S, Lee SB: **Rare codon clusters at 5'-end influence heterologous expression of archaeal gene in *Escherichia coli*.** *Protein Expr Purif* 2006, **50**:49-57.
35. Koma D, Sawai T, Harayama S, Kino K: **Overexpression of the genes from thermophiles in *Escherichia coli* by high-temperature cultivation.** *Appl Microbiol Biotechnol* 2006, **73**:172-180.
36. Albers S-V, Driessen AJ: **Conditions for gene disruption by homologous recombination of exogenous DNA into the *Sulfolobus solfataricus* genome.** *Archaea* 2007, **2**:145-149.
37. Worthington P, Hoang V, Perez-Pomares F, Blum P: **Targeted disruption of the alpha-amylase gene in the hyperthermophilic archaeon *Sulfolobus solfataricus*.** *J Bacteriol* 2003, **185**:482-488.
38. Martin A, Yeats S, Janekovic D, Reiter WD, Aicher W, Zillig W: **SAV 1, a temperate u.v.-inducible DNA virus-like particle from the archaeobacterium *Sulfolobus acidocaldarius* isolate B12.** *EMBO J* 1984, **3**:2165-2168.
39. Clore AJ, Stedman KM: **The SSV1 viral integrase is not essential.** *Virology* 2007, **361**:103-111.
40. Aucelli T, Contursi P, Girifoglio M, Rossi M, Cannio R: **A spreadable, non-integrative and high copy number shuttle vector for *Sulfolobus solfataricus* based on the genetic element pSSVx from *Sulfolobus islandicus*.** *Nucleic Acids Res* 2006, **34**:e114.
41. Berkner S, Grogan D, Albers SV, Lipps G: **Small multicopy, non-integrative shuttle vectors based on the plasmid pRN1 for *Sulfolobus acidocaldarius* and *Sulfolobus solfataricus*, model organisms of the (cren)-archaea.** *Nucleic Acids Res* 2007, **35**:e88.
42. Albers SV, Jonuscheit M, Dinkelaker S, Ulrich T, Kletzin A, Tampe R, Driessen AJ, Schleper C: **Production of recombinant and tagged proteins in the hyperthermophilic archaeon *Sulfolobus solfataricus*.** *Appl Environ Microbiol* 2006, **72**:102-111.
43. Schelert J, Dixit V, Hoang V, Simbahan J, Drozda M, Blum P: **Occurrence and characterization of mercury resistance in the hyperthermophilic archaeon *Sulfolobus solfataricus* by use of gene disruption.** *J Bacteriol* 2004, **186**:427-437.
44. Du Plessis CA, Batty JD, Dew DW: **Commercial applications of thermophile bioleaching.** In *Biomining*. Edited by Rawlings DE, Johnson DB. Springer-Verlag; 2007.
45. Mikkelsen D, Kappler U, McEwan AG, Sly LI: **Archaeal diversity in two thermophilic chalcopyrite bioleaching reactors.** *Environ Microbiol* 2006, **8**:2050-2056.
46. Rawlings DE, Johnson DB: **The microbiology of biomining: development and optimization of mineral-oxidizing microbial consortia.** *Microbiology* 2007, **153**:315-324.
- Consortia optimization strategy considerations based on microbiology of biocatalysts, as well as chemical engineering characteristics of biomining operations.
47. Chen L, Brugger K, Skovgaard M, Redder P, She Q, Torarinsson E, Greve B, Awayez M, Zibat A, Klenk HP et al.: **The genome of *Sulfolobus acidocaldarius*, a model organism of the Crenarchaeota.** *J Bacteriol* 2005, **187**:4992-4999.
48. Kawarabayasi Y, Hino Y, Horikawa H, Jin-no K, Takahashi M, Sekine M, Baba S, Ankaï A, Kosugi H, Hosoyama A et al.: **Complete genome sequence of an aerobic thermoacidophilic crenarchaeon, *Sulfolobus tokodaii* strain 7.** *DNA Res* 2001, **8**:123-140.
49. She Q, Singh RK, Confalonieri F, Zivanovic Y, Allard G, Awayez MJ, Chan-Weiher CC-Y, Clausen Ib G, Curtis BA, De Moors A et al.: **The complete genome of the crenarchaeon *Sulfolobus solfataricus* P2.** *Proc Natl Acad Sci U S A* 2001, **98**:7835-7840.
50. Auernik KS, Maezato Y, Blum PH, Kelly RM: **The genome sequence of the metal-mobilizing, extremely thermoacidophilic archaeon *Metallosphaera sedula* provides insights into bioleaching-associated metabolism.** *Appl Environ Microbiol* 2008, **74**:682-692.
- This study discusses genome features relevant to biomining and genome-wide transcriptional response to the presence of ferrous iron.
51. Castresana J, Lubben M, Saraste M: **New archaeobacterial genes coding for redox proteins: implications for the evolution of aerobic metabolism.** *J Mol Biol* 1995, **250**:202-210.
52. Lubben M, Arnaud S, Castresana J, Warne A, Albracht SP, Saraste M: **A second terminal oxidase in *Sulfolobus acidocaldarius*.** *Eur J Biochem* 1994, **224**:151-159.

53. Purschke WG, Schmidt CL, Petersen A, Schafer G: **The terminal quinol oxidase of the hyperthermophilic archaeon *Acidianus ambivalens* exhibits a novel subunit structure and gene organization.** *J Bacteriol* 1997, **179**:1344-1353.
54. Hiller A, Henninger T, Schafer G, Schmidt CL: **New genes encoding subunits of a cytochrome bc₁-analogous complex in the respiratory chain of the hyperthermoacidophilic crenarchaeon *Sulfolobus acidocaldarius*.** *J Bioenerg Biomembr* 2003, **35**:121-131.
55. Kappler U, Sly LI, McEwan AG: **Respiratory gene clusters of *Metallosphaera sedula* – differential expression and transcriptional organization.** *Microbiology* 2005, **151**:35-43.
56. Bathe S, Norris PR: **Ferrous iron- and sulfur-induced genes in *Sulfolobus metallicus*.** *Appl Environ Microbiol* 2007, **73**:2491-2497.
- Identification of a new terminal oxidase cluster and ETC components involved in ferrous iron oxidation in selected *Sulfolobus* species.
57. Urich T, Gomes CM, Kletzin A, Frazao C: **X-ray structure of a self-compartmentalizing sulfur cycle metalloenzyme.** *Science* 2006, **311**:996-1000.
- This study describes structural resolution of an intracellular sulfur oxidoreductase and the implications for its mechanism of operation via creation of a microenvironment.
58. Kletzin A: **Metabolism of inorganic sulfur compounds in archaea.** In *Archaea: Evolution, Physiology, and Molecular Biology*. Edited by Garrett RA, Klenk HP. Blackwell; 2007:261-276.
- A thorough summary of known inorganic sulfur metabolism in archaea and comparison to bacterial metabolism.
59. Berg IA, Kockelkorn D, Buckel W, Fuchs G: **A 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon dioxide assimilation pathway in Archaea.** *Science* 2007, **318**:1782-1786.
- Outline of a new inorganic carbon fixation cycle in archaea and phylogenetic diversity of microorganisms containing key enzymes.
60. Hugler M, Krieger RS, Jahn M, Fuchs G: **Characterization of acetyl-CoA/propionyl-CoA carboxylase in *Metallosphaera sedula*. Carboxylating enzyme in the 3-hydroxypropionate cycle for autotrophic carbon fixation.** *Eur J Biochem* 2003, **270**:736-744.
61. Alber B, Olinger M, Rieder A, Kockelkorn D, Jobst B, Hugler M, Fuchs G: **Malonyl-coenzyme A reductase in the modified 3-hydroxypropionate cycle for autotrophic carbon fixation in archaeal *Metallosphaera* and *Sulfolobus* spp.** *J Bacteriol* 2006, **188**:8551-8559.
62. Alber BE, Kung JW, Fuchs G: **3-Hydroxypropionyl-coenzyme A synthetase from *Metallosphaera sedula*, an enzyme involved in autotrophic CO₂ fixation.** *J Bacteriol* 2008, **190**:1383-1389.
63. Tang TH, Polacek N, Zywicki M, Huber H, Brugger K, Garrett R, Bachellerie JP, Huttenhofer A: **Identification of novel non-coding RNAs as potential antisense regulators in the archaeon *Sulfolobus solfataricus*.** *Mol Microbiol* 2005, **55**:469-481.
64. Omer AD, Zago M, Chang A, Dennis PP: **Probing the structure and function of an archaeal C/D-box methylation guide sRNA.** *RNA* 2006, **12**:1708-1720.
65. Ciammaruconi A, Gorini S, Londei P: **A bifunctional archaeal protein that is a component of 30S ribosomal subunits and interacts with C/D box small RNAs.** *Archaea* 2007, **2**:151-158.
66. Lillestol RK, Redder P, Garrett RA, Brugger K: **A putative viral defence mechanism in archaeal cells.** *Archaea* 2006, **2**:59-72.
67. Sorek R, Kunin V, Hugenholtz P: **CRISPR – a widespread system that provides acquired resistance against phages in bacterial and archaea.** *Nat Rev Microbiol* 2008, **6**(3):181-186.
68. Sleytr UB, Egelseer EM, Ilk N, Pum D, Schuster B: **S-layers as a basic building block in a molecular construction kit.** *FEBS J* 2007, **274**:323-334.
69. Engelhardt H: **Are S-layers exoskeletons? The basic function of protein surface layers revisited.** *J Struct Biol* 2007, **160**:115-124.
70. Engelhardt H: **Mechanism of osmoprotection by archaeal S-layers: a theoretical study.** *J Struct Biol* 2007, **160**:190-199.
71. Bode ML, Buddoo SR, Minnaar SH, du Plessis CA: **Extraction, isolation, and NMR data of the tetraether lipid calditoglycerocaldarchaeol (GDNT) from *Sulfolobus metallicus* harvested from a bioleaching reactor.** *Chem Phys Lipids* 2008, **154**(2):94-104.
72. Huber H, Stetter KO: **Desulfurococcales.** In *The Prokaryotes*, edn 3. Edited by Dworkin M, Falkow S, Rosenberg E, Schleifer K, Stackebrandt E. Springer; 2006:52-68.
73. Huber H, Huber R, Stetter KO: **Thermoproteales.** In *The Prokaryotes*, edn 3. Edited by Dworkin M, Falkow S, Rosenberg E, Schleifer K, Stackebrandt E. Springer; 2006:10-22.